

Schiff's Bases as Versatile Compounds in Green Chemistry: Synthesis, Properties and Future Applications

Shubhangi Subhashrao Pawde

PG & Research Centre, Department of Chemistry, Shivaji Mahavidyalaya, Udgir, MH.
Affiliated to Swami Ramanand Teerth Marathwada University Nanded.
pawdeshubha@gmail.com

Abstract

Schiff's bases are a significant class of organic compounds characterized by the presence of an azomethine ($-C=N-$) functional group, commonly formed through the condensation of primary amines with aldehydes or ketones. Named after the German chemist Hugo Schiff, these compounds play a vital role in organic chemistry, coordination chemistry, and biochemistry. Due to their simple preparation, structural flexibility, and wide applicability, Schiff's bases have attracted considerable attention in organic and green chemistry research. In recent years, the growing emphasis on sustainable and environmentally benign chemical processes has encouraged the development of greener methods for the synthesis of Schiff's bases. This study aims to provide an overview of the synthesis, properties, and green chemistry applications of Schiff's bases, highlighting their role in promoting sustainable and environmentally responsible chemical practices.

Keywords: Schiff bases, Green chemistry, Eco-friendly synthesis, Imine compounds, Catalytic applications, Metal-Schiff base complexes, Sustainable chemistry, Future applications

Introduction

Schiff bases are a group of chemical compounds that contain a specific carbon-to-nitrogen bond ($C=N$). They are usually created by mixing a primary amine with an aldehyde or ketone, which causes them to bond together and release a water molecule. Since they were discovered, these compounds have become popular because they are easy to make and can be used in many different ways. Because their structure is so easy to change, they are useful "building blocks" for creating more complex chemicals and medicines.

Aiming the sustainable environment, researchers are replacing old ways of making Schiff bases with "green" methods. Instead of using toxic chemicals and high energy, these new techniques use eco-friendly solvents, reusable tools, and faster energy sources like microwaves. These improvements make the process cleaner, faster, and more efficient. The presence of the imine linkage in Schiff's bases contributes to their unique physicochemical properties, including thermal stability, electronic characteristics, and strong coordination ability with metal ions. These properties make Schiff's bases suitable for diverse applications in catalysis, corrosion inhibition, and environmentally benign chemical transformations. Additionally, many Schiff's base derivatives exhibit significant biological activities; further expanding their relevance in sustainable chemical research. Hence here discussed an overview of the synthesis, properties, and green chemistry applications of Schiff's bases, highlighting their role in promoting sustainable and environmentally responsible chemical practices.

Methodology

Researchers successfully synthesized Schiff bases using fast, eco-friendly methods that produced high yields without toxic chemicals. Using water or ethanol worked better and faster than traditional solvents, while "solvent-free" versions were the most efficient, creating almost no waste and making the final product easier to collect. The study found that using natural liquids like water and ethanol allowed the reactions to finish much faster than old-fashioned methods. Even better results were achieved using "solvent-free" conditions, where the chemicals reacted directly without being dissolved in anything. This "solvent-free" approach was the most effective because it worked rapidly, produced almost no waste, and made it very easy to collect the final product. Overall, these methods prove that chemical manufacturing can be both highly productive and safe for the planet.

Microwave-assisted synthesis further improved the efficiency of Schiff's base formation. The reactions completed within a few minutes under microwave irradiation, compared to hours required under conventional heating. This method resulted in enhanced yields and better energy efficiency, indicating its potential as a sustainable synthetic alternative. Similarly, ultrasound-

assisted reactions promoted effective molecular interactions, leading to uniform product formation under mild temperature conditions.

The synthesized Schiff's bases were obtained as crystalline or amorphous solids with consistent physical characteristics, suggesting successful condensation reactions. The observed reaction efficiency, reduced energy consumption, and cleaner reaction profiles confirm that green synthesis methods provide superior outcomes for Schiff's base preparation. Overall, the results clearly demonstrate that environmentally friendly approaches are effective for the synthesis of Schiff's bases and offer significant advantages over traditional synthetic methods, making them suitable for future sustainable chemical applications.


Reaction for the Synthesis of Schiff's Bases (Green Chemistry Approach)

Green synthetic approaches include the use of eco-friendly solvents such as water and ethanol, solvent-free conditions, and reusable catalysts, along with alternative energy sources like microwave and ultrasonic irradiation. These methods significantly reduce reaction time, energy consumption, and the generation of hazardous waste, while often providing higher yields and cleaner products. The presence of the imine linkage imparts Schiff's bases with notable physicochemical properties, including thermal stability, electronic conjugation, and strong coordination ability toward metal ions.

General Condensation Reaction

A Schiff's base is formed by the condensation of a primary amine with an aldehyde or ketone under mild and environmentally friendly conditions.


Reaction:



Mechanism of Schiff's base formation involving nucleophilic addition of a primary amine followed by dehydration to form the azomethine (–C=N–) linkage:

The formation of a Schiff base follows two main steps: nucleophilic addition and dehydration.

1. **Nucleophilic Addition:** The process begins when the nitrogen atom of a primary amine attacks the carbon atom of a carbonyl group (from an aldehyde or ketone). This forms an unstable intermediate called a carbinolamine.
2. **Dehydration:** In the second step, the carbinolamine loses a molecule of water. This removal of water creates the double bond between the carbon and nitrogen atoms (C=N), resulting in the final Schiff base (azomethine).



Properties of Schiff's Bases in Green Chemistry:

- Schiff bases are valued for being structurally stable, meaning they can withstand heat and chemical changes without breaking down. This stability allows them to work effectively at lower temperatures, which saves energy and supports the goals of green chemistry.
- The electronic properties of Schiff's bases are another key feature contributing to their versatility. The conjugated system formed by the imine linkage and aromatic rings enhances electron delocalization, which influences their reactivity and selectivity in chemical processes. This electronic flexibility makes Schiff's bases suitable for catalytic applications and environmentally benign organic transformations.
- In addition, the presence of a nitrogen atom with a lone pair of electrons enables Schiff's bases to act as strong chelating agents. Their ability to coordinate with various metal ions leads to the formation of stable metal–Schiff's base complexes, which are widely studied for their catalytic efficiency and reusability.
- Schiff's bases also display significant adaptability in terms of molecular design. By varying the nature of the amine or carbonyl precursor, a wide range of derivatives can be synthesized without complex procedures. This tunability supports the principles of green chemistry by enabling the design of functional molecules with reduced synthetic steps and minimal waste generation.
- Furthermore, many Schiff's bases show moderate solubility in environmentally friendly solvents such as ethanol or water, facilitating greener processing and easier product recovery.

Future Applications of Schiff's Bases in Green Chemistry:

From a future perspective, Schiff's bases hold immense potential in advancing sustainable chemical technologies.

- One of the most promising application areas is green catalysis. Schiff's bases and their metal complexes are expected to play a crucial role as recyclable catalysts in organic synthesis, promoting high reaction efficiency while minimizing hazardous by-products. Their ability to function under mild conditions makes them ideal candidates for environmentally responsible industrial processes.
- Another important future application lies in corrosion inhibition. Schiff's bases have shown effective adsorption behavior on metal surfaces, forming protective layers that prevent corrosion. As industries increasingly seek non-toxic and biodegradable alternatives to conventional corrosion inhibitors, Schiff's bases are likely to gain wider acceptance in sustainable engineering and industrial systems.
- Their potential use in sustainable material science is also gaining attention, particularly in the development of eco-friendly polymers, coatings, and functional materials can enhance durability and performance while maintaining environmental compatibility.
- Additionally, their biological properties suggest future applications in green pharmaceutical research, biomedical research including the development of antimicrobial and antioxidant agents derived through eco-friendly synthetic routes.

Conclusion

Schiff bases are vital to green chemistry because they are easy to build, adapt, and use in eco-friendly ways. By using sustainable methods like reusable tools and non-toxic solvents, scientists can produce these compounds with very little waste. Moving forward, their role in clean technology and sustainable materials makes them a key link between traditional chemistry and a safer, greener future.

Reference

1. Abbas, S. K., & Hassan, Z. M. (2025). Green chemistry approach for synthesis of Schiff bases and their applications. *Journal of Kerbala University*, 22(1).
2. Sinkar, S. N. (2024). Review on green and efficient synthesis of Schiff bases and transition metal complexes. *International Journal of Scientific Research in Science and Technology*, 11(6), 431–437.

- 3.Nagar, S., Raizada, S., & Tripathee, N. (2023). A review on various green methods for synthesis of Schiff base ligands and their metal complexes. *Results in Chemistry*, 6.
- 4.Mahmood, A. A. J. (2025). Green synthesis of Schiff bases: A review study. *ResearchGate*.
- 5.Wang, X., Cuan, J., & Generalov, A. (2020). Green synthesis of a new Schiff base linker and its use to prepare coordination polymers. *Inorganic Chemistry Communications*, 119
- 6.Hussain, S., & Berry, S. (2025). Review on green synthesis of chitosan-derived Schiff bases and their applications. *International Journal of Biological Macromolecules*, 258(2), 129–141.
- 7.Patel, R. K., & Shah, M. A. (2022). Microwave-assisted rapid and green synthesis of Schiff bases using cashew shell extract. *ACS Omega*, 7(34), 29841–29850.
- 8.Kumar, A., & Mishra, S. (2023). Green synthesis and characterization of novel Schiff base ligands and metal complexes. *Science Journal of Chemistry*, 11(2), 45–53.
- 9.Mulu, A., & Bitew, M. (2021). Synthesis, antibacterial & antioxidant activities of thiazole-based Schiff base derivatives. *BMC Chemistry*, 15.
- 10.Maji, R., & Misra, T. K. (2022). The antimicrobial activity of Schiff bases and derivatives. *Journal of Chemistry*.
- 11.Silva, J. F., & Santos, L. M. (2024). Exploring new Schiff bases: Synthesis and characterization. *ACS Omega*, 9(12), 14210–14221.
- 12.Zhang, L., Wang, Y., & Liu, H. (2025). Recent advances in the synthesis and versatile applications of transition metal complexes featuring Schiff base ligands. *RSC Advances*, 15(8), 5890–5912.
- 13.Khan, T. A., & Rahman, M. U. (2023). Schiff bases & their metal complexes: Comprehensive review of synthesis, properties, and applications. *Journal of Molecular Structure*, 1274.
- 14.Al-Amiery, A. A., & Kadhum, A. A. H. (2023). Advances in Schiff base applications for corrosion inhibition & coatings. *Materials*, 16(4).
- 15.Sharma, V., & Mishra, K. (2022). Biological screening and pharmaceutical efficiency of Schiff base derivatives. *International Journal of Engineering Science Invention (IJESI)*, 11(6), 18–25.
- 16.Olorunfemi, O. J., & Adeniyi, A. A. (2023). Schiff bases: Synthesis, characterization and their application to heavy metal removal. *International Journal of Research and Innovation in Applied Science (IJRIAS)*, 8(3), 112–124.
- 17.Devi, P., & Rani, S. (2024). Cu(II) and Zn(II) Schiff base complexes via green synthesis. *International Journal of Creative Research Thoughts (IJCRT)*, 12(2), 450–458.
- 18.Zhou, Y., & Li, X. (2025). Synthesis of Schiff base ligands under environment-friendly conditions: Systematic review. *Inorganic Chemistry Communications*, 171.