

Structural Diversity of Schiff-Base Ligands and their Metal Complexes

S. R. Khandekar

Department Of Chemistry, Indira Mahavidyalaya, Kalamb (MS), India.
snehalkhandekar11@gmail.com

Abstract

Schiff-base ligands provide a modular platform for coordination chemistry because their imine functionality can be incorporated into diverse donor sets, denticities, substitution patterns and molecular architectures. Coordination with metal ions expands this diversity through changes in geometry, nuclearity, electronic structure and supramolecular organization. This review examines these structural variables from a structure property perspective, emphasizing how ligand architecture, donor-atom arrangement, rigidity, flexibility, metal identity, oxidation state and coordination number influence molecular and functional behaviour. Particular attention is given to electronic, photophysical, magnetic, catalytic and biological properties and to the complementary roles of crystallography, spectroscopy and computational chemistry in establishing reliable correlations. Although a large number of Schiff-base complexes have been reported, comparison across ligand families and metal centres remains difficult because apparently small structural changes can alter coordination modes, nuclearity and functional response. A multidimensional framework linking ligand architecture, metal binding, coordination geometry, electronic structure, intermolecular interactions and function is therefore proposed as a basis for more rational Schiff-base design.

Keywords: Schiff bases; metal complexes; coordination geometry; ligand architecture; structure–property relationships; photophysical properties; catalysis; bioinorganic chemistry

1. Introduction

Molecular design in coordination chemistry depends on controlling the spatial arrangement and electronic environment of a metal centre through ligand selection. Schiff bases are especially useful because their imine or azomethine group is readily formed by condensation of a carbonyl compound with a primary amine and can be incorporated into mono-, bi-, tri- and polydentate architectures. Salicylaldimines, salen- and salophen-type ligands, hydrazones, thiosemicarbazones and related systems consequently occupy a broad region of coordination chemistry. Their appeal arises not only from synthetic accessibility but from the ability to alter donor atoms, chelate-ring size, rigidity, conjugation, steric environment and protonation state while retaining a recognizable ligand framework [1–4].

The central scientific question is how this molecular diversity propagates through metal binding to determine coordination geometry, nuclearity, electronic structure and ultimately function. A change in the carbonyl precursor can modify steric demand and conjugation; changing the amine can alter denticity, flexibility and donor arrangement; and substitution can tune both electronic density and intermolecular interactions. These variables are coupled. A substituent introduced to strengthen metal binding may also change ligand conformation, while increased rigidity may stabilize one coordination mode but prevent the geometry preferred by another metal. Consequently, Schiff-base chemistry provides an unusually useful test bed for structure–property relationships, but the same diversity that permits property tuning also complicates generalization.

Recent structural work illustrates this point. Crystallographic and computational investigation of a fluorinated Schiff-base copper system revealed a dinuclear complex with distorted square-pyramidal copper environments and significant intra- and intermolecular interactions [5]. Zinc(II) Schiff-base complexes likewise demonstrate how hydrogen- and halogen-bonding interactions can influence the supramolecular environment beyond the primary coordination sphere [6]. Such observations show that a formal ligand formula is not sufficient to describe a complex: coordination geometry, nuclearity and solid-state organization may all change the resulting behaviour.

Schiff-base metal complexes have consequently been investigated in catalysis, sensing, bioinorganic chemistry, magnetism and photofunctional materials [2,7–10]. However, much of the literature is organized by application or metal ion, making it difficult to identify structural principles that remain valid across compound families. This review therefore adopts a structure-centred perspective, moving from ligand architecture to metal-induced structural diversity, then

Copyright: © by the Author(s)
This is an open access article distributed under the terms of the Creative Commons Attribution License and permitted unrestricted use, distribution, and reproduction in any medium.

to structure-dependent properties, characterization, applications and remaining gaps. The objective is to identify recurring design variables while recognizing where the chemistry remains system-specific.

2. Structural diversity of schiff-base ligands

The structural space of Schiff bases begins with the choice of carbonyl and amine precursors. Aromatic aldehydes can introduce rigid conjugated frameworks, whereas ketones and substituted carbonyl compounds provide additional steric and electronic control. Diamines and multifunctional amines generate multidentate ligands capable of imposing defined coordination environments. This modularity explains the large family of salen-, salophen-, hydrazone-, thiosemicarbazone- and related systems [1,3,4].

Denticity is one of the strongest structural controls. Monodentate ligands leave substantial freedom at the metal, while bidentate and higher-dentate ligands progressively constrain coordination. Yet denticity should not be equated with rigidity. Flexible multidentate ligands can adapt to different metals and produce several binding modes, whereas rigid ligands may impose a narrow geometry but develop strain when the metal preference is incompatible with the ligand pocket. The balance between preorganization and adaptability therefore becomes a key design variable.

Donor-atom identity further changes coordination behaviour. N,O donors are particularly common, while sulfur-containing Schiff bases provide softer, more polarizable donor environments. HSAB concepts are useful for anticipating metal preference, but actual structures also depend on oxidation state, steric effects, solvent, protonation and competing donor groups [3,11]. Substituents can alter imine basicity, phenolic acidity, metal–ligand covalency and charge-transfer energetics. Ortho substitution additionally changes steric accessibility and ring torsion, whereas remote substituents can modify electronic communication with smaller geometric effects.

Tautomerism and proton transfer add another dimension. Salicylaldimine and hydrazone systems may access alternative tautomeric forms, and metal coordination can shift these equilibria by stabilizing a deprotonated donor set. Structural studies of ONO pyridyl hydrazone complexes demonstrate that tautomerism, coordination and molecular structure can be closely coupled [12]. Thus, the C=N group should not be treated as an isolated functional group; its chemical role depends on the surrounding donor framework and protonation state.

Rigidity, planarity and conjugation also control intermolecular organization. Extended aromatic surfaces can promote π – π stacking, while phenolic, amide and hydrazone groups can create hydrogen-bond networks. These interactions may influence luminescence, recognition, magnetic exchange and crystal packing. A meaningful structure–property analysis must therefore consider both the first coordination sphere and the larger supramolecular environment.

3. Metal-induced structural diversity

Metal coordination introduces a second major source of structural variation. Ionic radius, oxidation state, electron configuration, coordination number and ligand-field preferences determine which coordination environments are accessible. Cu(II) often exhibits distorted tetragonal environments, Ni(II) can support square-planar or octahedral structures, Pd(II) and Pt(II) commonly favour square-planar geometry, while Zn(II) often accommodates tetrahedral or higher-coordinate arrangements. Fe, Co and Mn systems show additional variation because oxidation and spin state can change geometry and magnetic behaviour. Lanthanide ions form a distinct class in which large ionic radii permit high and variable coordination numbers [1,3,7].

The metal can also change ligand protonation and nuclearity. Deprotonated phenolate or carboxylate groups may bridge two or more metal ions, producing dinuclear, polynuclear or polymeric structures. Auxiliary ligands and counter-ions may become part of the coordination sphere, so the final complex can differ substantially from the simple ligand–metal stoichiometry anticipated from the synthesis. Nuclearity is especially important because metal–metal interactions and bridging pathways can generate cooperative magnetic, catalytic or energy-transfer behaviour.

High-nuclearity Schiff-base assemblies demonstrate the upper end of this structural spectrum. Recent Tb and Dy systems containing Schiff-base macrocycles have shown stepwise metal-templated assembly and functional behaviour including fluorescence sensing and single-molecule-magnet characteristics [13]. Such systems reinforce the idea that metal coordination can act as a construction principle for higher-order molecular architectures rather than simply producing a discrete coordination compound.

4. From molecular structure to function

The functional properties of Schiff-base complexes arise from the combined electronic and geometric environment created by the ligand and metal. Optical behaviour may involve ligand-centred, d–d, ligand-to-metal charge-transfer or metal-to-ligand charge-transfer transitions. Donor strength, conjugation, coordination geometry and metal–ligand covalency can therefore shift absorption and emission energies. Small structural changes may also modify nonradiative relaxation through changes in rigidity or molecular motion.

Lanthanide complexes illustrate the importance of the complete coordination environment. Schiff-base ligands can act as antennae that absorb excitation and transfer energy to lanthanide-centred states. Efficient sensitization depends on ligand excited-state energies, metal–ligand distances, charge-transfer states, aggregation and temperature rather than on chromophore structure alone [14]. Recent Tb and Dy complexes combine luminescence with magnetic relaxation, demonstrating that one coordination framework can couple optical and magnetic functions [15].

Magnetic behaviour is similarly structure-dependent. Transition-metal complexes can exhibit spin-state changes, anisotropy and exchange interactions, while lanthanide complexes can display slow magnetic relaxation when the coordination environment provides suitable anisotropy. In multinuclear systems, the identity and geometry of bridging ligands become particularly important because they control magnetic exchange pathways.

Catalysis provides another direct structure–function relationship. Schiff-base complexes have been investigated in oxidation, epoxidation, hydroformylation, cyclopropanation, C–C coupling, Michael addition and polymerization [8,9]. Catalytic behaviour depends on metal identity, oxidation state, open coordination sites, ligand donor strength and steric accessibility. A strongly chelating ligand can improve stability but may hinder substrate binding; a flexible ligand may facilitate coordination changes but can also permit decomposition. Thus, activity cannot be assigned to the ligand or metal independently.

Biological properties likewise depend on coordination-induced changes in charge, lipophilicity, redox behaviour and molecular geometry. Schiff-base complexes have been widely investigated for antimicrobial, antioxidant and anticancer effects [3,10,16]. However, comparisons across studies are limited by differences in assay conditions and biological models. Structural claims about activity are strongest when free ligand, metal salt and complex are compared under matched conditions.

Supramolecular organization provides an additional level of control. Hydrogen bonding, π -stacking, electrostatic interactions and halogen bonding can determine crystal packing and influence sensing, optical and magnetic behaviour. Recent zinc(II) Schiff-base structures demonstrate how secondary interactions can become an important part of the molecular architecture [6]. Structure–property relationships should therefore extend beyond metal–donor bonds to include the wider molecular and crystal environment.

5. Synthesis and structural control

Conventional Schiff-base synthesis is generally straightforward, but formation of the final metal complex is not always structurally deterministic. Solvent, pH, stoichiometry, temperature, counter-ion and crystallization conditions can influence protonation, coordination mode and nuclearity. Microwave, ultrasound, solvent-free and aqueous procedures have been developed to reduce reaction time and solvent use [17,18].

Synthetic methodology should therefore be evaluated not only by yield and reaction time but also by structural reproducibility. Microwave heating may change nucleation and thermal gradients, while sonochemical methods can alter mixing and local energy deposition. Green

protocols can reduce environmental impact, but a lower-solvent synthesis is not necessarily superior if it produces mixed phases or poorly reproducible coordination products. Recent reviews emphasize this balance between efficiency, sustainability and control [17,18].

6. Structural characterization and methodological challenges

Reliable structure–property relationships require complementary characterization. Single-crystal X-ray diffraction is the most direct method for defining molecular connectivity, coordination geometry, nuclearity and crystal packing when suitable crystals are available. Powder X-ray diffraction is valuable for phase identification but generally cannot replace a complete single-crystal structure for complex architectures.

FTIR can reveal changes in imine and donor-group vibrations, NMR is useful for diamagnetic species, UV–visible spectroscopy probes electronic transitions, and photoluminescence measurements reveal emissive states. EPR is particularly informative for paramagnetic centres such as Cu(II) and Mn(II), while electrochemistry can distinguish accessible redox processes. Thermal analysis provides information on solvent loss and decomposition. Mass spectrometry can support molecular composition, although gas-phase ions should not automatically be equated with solid-state structures.

DFT and related computational methods increasingly complement these measurements by examining molecular orbitals, charge distribution, geometry and excited states. Hirshfeld-surface and quantum-theory-of-atoms-in-molecules analyses can further clarify intermolecular interactions [5]. The strongest studies therefore combine structural, spectroscopic and computational evidence rather than assigning complete molecular structure from a single technique.

7. Structure–property relationships and knowledge gaps

Several broad trends are well established: increasing denticity generally increases geometric constraint; donor strength influences metal binding and electronic structure; conjugation affects optical behaviour; and metal identity strongly influences geometry, redox and magnetic properties. Yet these relationships are not universally transferable. A ligand modification that strengthens binding to one metal may alter tautomerism or steric congestion with another, while increased rigidity may improve photophysical stability but prevent a preferred coordination geometry.

The literature also contains a persistent separation between molecular structure and functional testing. Crystallographic studies may define a precise solid-state structure without establishing solution behaviour; photophysical studies may report strong emission without fully resolving the emissive species; catalytic studies may optimize yield without proving catalyst integrity; and biological studies may report activity without resolving the separate contributions of ligand and metal. These differences complicate direct comparison.

The principal knowledge gap is therefore not a shortage of Schiff-base structures but a shortage of broadly transferable, experimentally validated structure–property relationships. Three levels require stronger integration: molecular coordination structure, supramolecular organization and functional performance. Future studies should systematically vary one structural descriptor at a time, characterize both solution and solid states where relevant, and use standardized functional measurements. Data-driven methods may eventually assist prediction, but their usefulness will depend on consistent structural assignments and comparable experimental datasets.

8. Emerging applications

Schiff-base complexes continue to support diverse applications. In catalysis, salen- and related frameworks provide tunable coordination environments for selective transformations [8,9]. Immobilization on magnetic supports can facilitate separation and recycling, although support attachment may change accessibility and coordination geometry [19].

In bioinorganic and medicinal chemistry, metal coordination can modify charge, lipophilicity, redox properties and biomolecular interactions. Recent reviews cover antimicrobial, antioxidant and anticancer applications, including efforts to address drug resistance [10,16,20]. These

directions are scientifically promising, but practical development requires systematic toxicity, selectivity, stability and mechanistic studies.

Sensing and photofunctional materials exploit the optical and electronic consequences of ligand–metal interaction. Conjugated Schiff bases can respond to analytes through changes in fluorescence, absorption or proton-transfer equilibria. Lanthanide systems add characteristic emission, while multinuclear architectures can provide cooperative responses [13,14]. Metal Schiff-base materials have also been explored in optoelectronic and energy-related applications because conjugated ligands can combine optical properties with metal-centred redox chemistry [21]. In each case, molecular performance should be distinguished from complete device or application performance.

9. Conceptual framework for rational design

Figure 1 shows a useful framework for Schiff-base design is:




Figure 1. Structure–property–function relationship of Schiff-base metal complexes.

This sequence is not strictly linear. Metal binding can alter ligand protonation, while crystal packing can influence conformation and photophysics. Nevertheless, it provides a practical hierarchy for testing proposed structure–property relationships. Ligand descriptors include denticity, donor type, chelate-ring size, rigidity, flexibility, conjugation and substituent electronics. Metal descriptors include oxidation state, ionic radius, d-electron configuration and preferred coordination number. Supramolecular descriptors include hydrogen bonding, π -stacking, electrostatic contacts and packing density.

The field can therefore move toward rational design by combining controlled ligand series with crystallography, spectroscopy, computation and functional testing. The goal should be not simply to maximize one measured property, but to understand which structural features produce reproducible performance across related systems.

10. Scientific and practical significance

Understanding Schiff-base structural diversity is important because it connects fundamental coordination chemistry with molecular design. At the molecular level, it clarifies how donor arrangement, metal identity and geometry control electronic structure and reactivity. In catalysis, such understanding can support deliberate tuning of activity and selectivity. In sensing and photofunctional materials, it can connect conjugation and coordination with optical responses. In bioinorganic chemistry, it can help separate ligand, metal and coordination-induced contributions to biological behaviour.

The practical objective is to replace isolated structure or activity reports with reproducible design principles. This requires attention to stability, scalability, reproducibility and realistic operating conditions. The field will benefit most from studies that connect these requirements rather than optimizing a single property in isolation.

11. Conclusions

Schiff-base ligands provide an unusually adaptable platform in which changes in molecular architecture generate broad variation in metal coordination geometry, nuclearity, electronic structure and supramolecular organization. This diversity supports applications in catalysis, sensing, photophysics, magnetism and bioinorganic chemistry, but it also makes generalized structure–property relationships difficult to establish. Future progress should emphasize systematically varied ligand series, cross-metal comparisons, rigorous structural characterization, solution/solid-state correlation and standardized functional measurements. Computational and data-driven methods may become increasingly valuable once sufficiently consistent structural and property datasets are available. Ultimately, Schiff-base complexes should be treated as integrated molecular systems in which ligand architecture, metal identity, coordination geometry, electronic structure and supramolecular organization jointly determine function.

References

1. Ghanghas, P., Choudhary, A., Kumar, D., & Poonia, K. (2021). Coordination metal complexes with Schiff bases: Useful pharmacophores with comprehensive biological applications. *Inorganic Chemistry Communications*, 133, 108710.
2. Abu-Dief, A. M., & Mohamed, I. M. A. (2015). A review on versatile applications of transition metal complexes incorporating Schiff bases. *Beni-Suef University Journal of Basic and Applied Sciences*, 4, 119–133.
3. Schiff bases and their metal complexes: A review on the history, synthesis, and applications. (2023). *Inorganic Chemistry Communications*, 150, 110451.
4. Juyal, V. K., Pathak, A., Panwar, M., Thakuri, S. C., Prakash, O., Agrwal, A., & Nand, V. (2023). Schiff base metal complexes as a versatile catalyst: A review. *Journal of Organometallic Chemistry*, 999, 122825.
5. Structural Investigation of Schiff Base Ligand and Dinuclear Copper Complex: Synthesis, Crystal Structure, Computational, and Latent Fingerprint Analysis. (2024). *ACS Omega*.
6. Gishan, M., Middya, P., Drew, M. G. B., Frontera, A., & Chattopadhyay, S. (2024). Synthesis, structural characterization, and theoretical analysis of novel zinc(II) Schiff base complexes with halogen and hydrogen bonding interactions. *RSC Advances*.
7. Barreras-Contreras, et al. (2025). Schiff Bases and Metal Complexes as Multifunctional Platforms: Bridging Bioinorganic Chemistry, Catalysis, Sensing, and Energy Applications. *Bioinorganic Chemistry and Applications*, 2025, 8829417.
8. Rakhtshah, J. (2022). A comprehensive review on the synthesis, characterization, and catalytic application of transition-metal Schiff-base complexes immobilized on magnetic Fe₃O₄ nanoparticles. *Coordination Chemistry Reviews*, 467, 214614.
9. Metal complexes with ONS donor Schiff bases: A review. (2022). *Polyhedron*, 217, 115692.
10. Recent advances in biological and medicinal profile of Schiff bases and their metal complexes: An updated version (2018–2023). (2024). *Results in Chemistry*, 7, 101350.
11. Verma, C., & Quraishi, M. A. (2021). Recent progresses in Schiff bases as aqueous phase corrosion inhibitors: Design and applications. *Coordination Chemistry Reviews*, 446, 214105.
12. Metal complexes incorporating tridentate ONO pyridyl hydrazone Schiff base ligands: Crystal structure, characterization and applications. (2023). *Coordination Chemistry Reviews*.
13. Li, H.-Y., et al. (2024). Two Octa-Nuclear {Ln₈} (Ln = Tb, Dy) Complexes with “4 + 4” Covalent Organic Macrocycles: Structure, Solution Behavior, Luminescence, and Magnetic Properties. *Crystal Growth & Design*.
14. Factors that influence the energy transfer pathways in luminescent lanthanide Schiff base complexes. (2024). *Inorganica Chimica Acta*, 572, 122324.
15. Gusev, A., Nemec, I., Herchel, R., Baluda, Y., Babeshkin, K., Efimov, N., Kiskin, M., & Linert, W. (2024). Lanthanide(III) SMMs with cationic and anionic complex fragments formed by a Schiff base: Structure, luminescence, magnetic properties and ab initio calculations. *Dalton Transactions*, 53, 11531–11542.

16. Podolski-Renić, A., et al. (2024). Schiff bases and their metal complexes to target and overcome (multidrug) resistance in cancer. *European Journal of Medicinal Chemistry*, 270, 116363.
17. A comprehensive review on the synthesis and applications of Schiff base ligand and metal complexes: A comparative study of conventional heating, microwave heating, and sonochemical methods. (2023). *Inorganic Chemistry Communications*.
18. Sustainable synthetic endeavors of pharmaceutically active Schiff bases and their metal complexes: A review on recent reports. (2024). *Tetrahedron*, 153, 133836.
19. Verma, C., & Quraishi, M. A. (2021). Recent progresses in Schiff bases as aqueous phase corrosion inhibitors: Design and applications. *Coordination Chemistry Reviews*, 446, 214105.
20. Advancement in the synthesis of metal complexes with special emphasis on Schiff base ligands and their important biological aspects. (2024). *Results in Chemistry*, 7, 101222.
21. Energy materials based on metal Schiff base complexes. (2018). *Coordination Chemistry Reviews*.