

MYCORRHIZAL STATUS AND ISOLATION OF SPORES FROM RHIZOSPHERIC SOIL OF VIGNA MUNGO IN ARDHAPUR REGION OF NANDED DISTRICT**S.B. Wankhede**Rajiv Gandhi Mahavidyalaya Mudkhed, Dist. Nanded, Maharashtra India
drsavitawankhede@gmail.com**ABSTRACT**

Arbuscular mycorrhizal fungi play an important role in the mobilization nutrients and enhancing plant growth. Mycorrhizal status and isolation of spores from rhizospheric soil of *Vigna mungo* in Ardhapur region of Nanded district. The rhizospheric soil was screened for spore density and population. The spore density were recorded as 280 spores per 100gm of soil and The spore population mainly consist of different species of Arbuscular mycorrhizal such as mainly consist of *Glomus*, *Acaulospora* and *Gigaspora*. spores were identified by using the manual of (Schenck and Perez, 1990). *Glomus fasciculatum* with subtending hyphae. Rounded shaped *Glomus reticulatum* and *Glomus* species. *Glomus fragilistatum*, *Glomus citricolla*, ruptured wall of *Glomus macrocarpum* and *Glomus globiform*, *Glomus mosseae* and *Acaulospora laevis*, *Acaulospora* sp. and *Scutellospora pellicida*. hyphae, vesicles, and Arbuscular seen in Whole mount of root was analyzed for the root colonization by using the method Phillips and Hymen (1970). The % colonization was 70 to 75% in root of *Vigna mungo*.

Keywords: Arbuscular Mycorrhizal fungi, Root colonization, *Vigna mungo*

Introduction

German Botanist Frank (1885) coined the term mycorrhizae for the first time to designate the symbiotic relationship between the fungi and plant roots. Since then scientists started exploiting them for the welfare of mankind. The term 'mycorrhiza' in its broadest sense is the non-pathogenic association of fungi and the roots of higher plants. The root- fungus association is symbiotic and the whole association is being considered as a "functionally distinct organ" involved in mineral nutrient uptake from the soil. (Kar, 1993). Mycorrhizal fungi are having intimate association with roots of higher plants forming a symbiotic relationship providing nutrients to the plants. The Arbuscular Mycorrhizal diversity in herbaceous vegetation medicinal plants, in halophytes plants have been investigated by many workers [Bagyaraj, D. J. (2014) Kannan, K. and Lakshminarashiman, C. (1988) Kumar., *et. al* (2013). Mulla, R. M *et. al.*, (1994) Mulani., R. M *et. al.*, (2004) Mulani, R. M and Waghmare, S. S. (2012). Mulani, R. M and Prabhu, R. R. (2002). Parameswaran, P and Augustine, B.(1988). Isolation and identification of arbuscular mycorrhizal fungi from agricultural fields of Vietnam investigated by (Sasvari *et.al.*, 2012). Growth and biomass of *Piper longum* L was

increased with inoculation of arbuscular mycorrhizal fungi. (Seema and Rajkumar,2015). Essential oil production, nutrient uptake and root colonization in basil was increased with inoculation arbuscular mycorrhizal fungi. (Mirhassan *et.al.*,2010). Black gram (*Vigna mungo* L.) also known as urd bean in India is one of the most important cultivated pulse crops of the 'Vigna' group and cultivation from ancient times. It is grown in various agro-ecological conditions and cropping systems with diverse agricultural practices, both in rainy (kharif) and post rainy (rabi) seasons . It is grown in summer and winter and accounts for 28 per cent of world grain legume production.(Ajaykumar;2022). *Vigna mungo* seeds are traditionally used as food and leaves as vegetable. Seeds are used as nervine tonic for the treatment of male sterility problems and act as a good aphrodisiac agent. It is also used to treat urinary reflex disorder. Oil of seeds is used to treat neurological problems like hemiplegia, polio myelitis and rheumatological problem.(Zaheer *et.al.*,2020).

Materials and Methods**Isolation of spores by using wet-sieving method. (Gerdman and Nicolson; 1963)**

Spore extraction is involved in three sub steps such as wet-sieving, sedimentation, flotation. Mix 5 gm of soil in 250 ml of luck warm water

in a beaker until all aggregates disperse to a uniform suspension. Allow the heavier particles to settle down. Filter the suspension through 710 μm sieve to remove large organic matter and roots. Then solution was sieved through series of sieves i.e. 710 μm , 210 μm , 150 μm , 75 μm , 45 μm and 25 μm respectively. Content of each sieves i.e. 210 μm , 150 μm , 75 μm , 45 μm and 25 μm was taken separately on blotting paper in petriplate and This petriplate was observed under stereo zoom binocular microscope.

Percentage of root colonization. (Phillips and Hayman, 1970).

Young root segments were taken in test tube adding 10% KOH and it autoclaved at 15 lbs

for 1 hr. After 10 minute 10% KOH was removed from test tube then root segments were washed under tap water with 2 to 3 times . Then 10 ml 1N HCL was added and were kept for 5 minute for neutralization of root tissue. Then HCL was removed and washed the root segments 2 to 3 times with tap water. After 30 minute root segments stained with cotton blue and kept for 24 hrs. After 24 hrs root segments mounted on slide with Acetic acid – glycerol (1:1v/v). Seal the corners of the cover slip with DPX, root colonization was observed under compound microscope. Then % of Arbuscular Myccorrhizal fungal colonization calculated by using this formula.

$$\text{Percent of mycorrhizal colonization} = \frac{\text{Number of root segments colonized}}{\text{Total number of root segments examined}} \times 100$$

Result and Discussion

The roots of *Vigna mungo* showed 70 to 75% Mycorrhizal colonization and the rounded, vesicles were prominent. The rhizospheric soil was screened for spore density and population. The spore density were recorded as 280 spores per 100gm of soil and The spore population mainly consist of different species of Arbuscular mycorrhizal such as mainly consist of *Glomus*, *Aculospora* and *Gigaspora*. spores were identified by using the manual of (Schenck and Perez, 1990). *Glomus fasciculatum* with subtending hyphae. Rounded shaped *Glomus reticulatum* and *Glomus species*. *Glomus fragilistatum*, *Glomus citricolla*, ruptured wall of *Glomus macrocarpum* and *Glomus globiformum*, *Glomus mosseae* and *Acaulospora laevis*, *Acaulospora sp.* and *Scutellospora pellicida*, *Scutellospora auriglobosa* and *Scutellospora calspora*. *Gigaspora rosea*. Similar observation made by Sasvari *et. al.*, (2012) in their studies highest number of spores found in the tomato and peanuts at agricultural field of Vietnam.

The roots of *Aloe vera* showed 90 % root colonization and spore density was recorded as 250 spores per 100 gm of soil. Such observation were made by Mulani and Waghmare, (2012). The presence of large number of spore with varied population of spores indicated their universal occurrence in

the soil of university campus. Such observations were made by Mulani and Prabhu. (2002), Mulani *et.al.*, (2004), Prabhu(2002) and Sathe (2005). Mulani and Prabhu had observed highest count of chlamydospores occurring in the root zone soil of *Dipcadi saxorum*. The murmy soil with moisture % and low humidity with high temperature fevers more chlymadospore formation. Similar observations were made by Harinikumar and Bagyaraj (1988) and Bagyaraj (1995) in tropical soil. Recently Pawar and Kakde (2012) have carried out the studies on the AMF associated with some medicinal plants from Mumbai region. They reported eight different species of *Glomus* namely *G. aggregatum*, *G. Boreale*, *G. fasciculatum*, *G. geosporum*, *G. heterosporum*, *G. segmentatum*, *G. tortuosum*, *G. radiatum* associated with the selected medicinal plants.

Root colonization of *Vigna mungo* showing in Fig : a,b c, d, e, f, g,i (Plate-I). Magnified view of rounded vesicles, Hyphae and Arbuscles seen in whole mount of root of *Vigna mungo* (40x, 100x). different spores were isolated from rhizospheric soil of *Vigna mungo* from Ardhapur region in Nanded District. fig-j: *Glomus fragilistatum*. fig-k: *Glomus citricolla*.fig-l: ruptured wall of *Glomus macrocarpum*. fig-m: *Glomus globiformum*.

PLATE-I





PLATE-II



References

1. Ajaykumar, P. Prabakaran, K. Sivasabari(2022). Growth and Yield Performance of Black Gram (*Vigna mungo* L.) under Malabar Neem(*Melia dubia*) Plantations in Western Zone of Tamil Nadu. *An International Journal*, Volume 45 Issue 2: 182-188.
2. Bagyaraj, D. J. (2014). Mycorrhizal fungi. *Proc Indian Natn Sci Acad.* 80(2): 415-428.
3. Bagyaraj, D. J. (1995). Influence of agricultural practices on vesicular arbuscular mycorrhizal fungi in soil. *Journal soil biol. Ecol.* 15(2):109-116.
4. Gerdman, J. W. and Nicolson, T. H. (1963). Spores of mycorrhizal *Endogone* species extracted from the soil by wet sieving and decanting. *Trans. Br. Mycol. Soc.* 46:235-244.
5. Kannan, K. and Lakshminarashiman, C. (1988). Survey of VAM of maritime strand plants of Po Calimere. In-First Asian conference on Mycorrhizae, C.A.S. in Botany, Madras. 29(31): 53-55.
6. Kumar, A. Chhavi, M. and Aggrawal, A. (2013). Biodiversity of Endophytic mycorrhizal fungi associated with some medicinal plants of Himachal Pradesh. *Asian J. of Adv. Basic sci.* 1 (1): 26-29.
7. Mulla, R. M. and Kanade, A. M. (1994). VAM Mycorrhizal colonization in grasses of Bombay. *J. Rayat Shikshan Sanstha Satara* : 56-65.

8. Mulani, R. M, Prabu, R. R. and Dinkaran, M. (2004). Occurrence of vesicular Arbuscular Mycorrhizae (VAM) in the roots of *Phyllanthus fraternus* Webster. *Mycorrhiza News*. 14 (2):11-14.
9. Mulani, R. M and Waghmare, S. S. (2012). Assessment of occurrence of Thermo tolerant Arbuscular Mycorrhizal Fungi in the Roots and Rhizospheric soil of *Aloe vera* (.) Burn.f. Online international journal interdisciplinary research journal. 2(4): 22-27.
10. Mulani, R. M. and Prabhu, R. R. (2002). A seasonal variation in Arbuscular Mycorrhizal (VAM) colonization in the roots of *Dipcadi saxorum* Blatt and chlamydospores in the rhizospheric soil from Mumbai. *J. sol. Biol. & ECOL*. 20 (172): 47-50.
11. MirHassan, R., Abbas. H., Mohsen B., Younes., R and Fatemeh., S.(2010) Effects of arbuscular mycorrhizal (AM)fungi on growth, essential oil production and nutrients uptake in basil. *Journal of Medicinal Plants Research* Vol.4(21), pp. 2222-2228.
12. Parameswaran, P and Augustine, B. (1988). Distribution and ecology of VAM in a scrub jungle. In-First Asia conference on Mycorrhizae, C. A. S. in Botany, Madras. 29(31): 91-99.
13. Pawar, J. S and Kakde, U. B (2012). Study of arbuscular Mycorrhiza associated with some important medicinal plants suburban area of Mumbai. Online international journal interdisciplinary research journal. II. 116-127.
14. Phillips, J. M. and Hayman, D. S. (1970). Improved procedure for clearing roots and staining parasitic and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. *Trans. Br. Mycol. Soc.* 55: 152-160.
15. Prabhu, R. R. (2002). Survey of soil of Mumbai and Adjoining areas for native VAM and their multiplication and effect of their inoculation on local crops as biofertilizers. A Ph.D thesis submitted to Mumbai University.
16. Sasavari, Z, Magurno. F, Galanics, D, Hang T. T, Hong Ha. T. T, Luyen .N. D, Huang .L and Posta .K (2012). Isolation and identification of Arbuscular Mycorrhizal Fungi from agricultural fields of Vietnam .*American Journal of plant sciences* 3, 1796-1801.
17. Sathe, V. D.(2005). Assessment of arbuscular mycorrhizal status in the soil on some forest plateaus of the Western ghats of Maharashtra. A Ph.D thesis submitted to the Mumbai University.
18. Seema, H. S and Rajkumar H. G. (2015) Effect of arbuscular mycorrhizal fungi on growth and biomass enhancement in *piper longum* L.(Piperaceae). *Int. J. Curr. Microbiol. App. Sci* (2015) 4(1): 11-18.
19. Zaheer. M, Ahamad,. S and Hasan, M(2020) Review of medicinal uses photochemistry and pharmacology of *Vigna mungo*(L). *Hepper Journal of Pharmacognosy and Phytochemistry*. 9(1): 1307-1309.